




Hi Carol,
You are nearing your surgery,
so now is a good time to start
preparing.

Please stop taking aspirin and
NSAIDs one week before surgery.

After surgery, you will need to
wear your sling for 4-6 weeks.
We will show you how to put it
on and take it off.



Every Step of the Way.



Keep your patients the center
of their surgical journey



EXAC coach™

Virtually coach your patients from surgical preparation through rehabilitation through the exacCoach™ patient text messaging service.



Reverse Total Shoulder Replacement
Anatomical Total Shoulder Replacement
Rotator Cuff Repair



Total Knee Replacement
Partial Knee Replacement
Revision Knee Replacement
Anterior Cruciate Ligament




Anterior Total Hip Replacement
Posterior Total Hip Replacement
Revision Hip Replacement

Key Benefits

Stay connected with your patients and caregivers through a fully customizable messaging system that automates personal text conversations, including words of encouragement, reminders, exercise recommendations and videos. Priced for simple integration into your practice and protocols.

Automated

Messages are activated based on surgery stage from preoperative care to rehabilitation.



Customizable

Fully personalized messages and schedule that best meet your practice's needs.

Timely

Orthopaedic surgeon-driven design helps anticipate appropriate cadence and timing of messages.



Post-op day 10

